

# Static Magnetic Field Therapy for Symptomatic Diabetic Neuropathy: A Randomized, Double-Blind, Placebo-Controlled Trial

Michael I. Weintraub, MD, FACP, FAAN, Gil I. Wolfe, MD, Richard A. Barohn, MD, Steven P. Cole, PhD, Gareth J. Parry, MD, Ghazala Hayat, MD, Jeffrey A. Cohen, MD, Jeffrey C. Page, DPM, Mark B. Bromberg, MD, Sherwyn L. Schwartz, MD, and the Magnetic Research Group

**ABSTRACT.** Weintraub MI, Wolfe GI, Barohn RA, Cole SP, Parry GJ, Hayat G, Cohen JA, Page JC, Bromberg MB, Schwartz SL, and the Magnetic Research Group. Static magnetic field therapy for symptomatic diabetic neuropathy: a randomized, double-blind, placebo-controlled trial. *Arch Phys Med Rehabil* 2003;84:736-46.

**Objective:** To determine if constant wearing of multipolar, static magnetic (450G) shoe insoles can reduce neuropathic pain and quality of life (QOL) scores in symptomatic diabetic peripheral neuropathy (DPN).

**Design:** Randomized, placebo-control, parallel study.

**Setting:** Forty-eight centers in 27 states.

**Participants:** Three hundred seventy-five subjects with DPN stage II or III were randomly assigned to wear constantly magnetized insoles for 4 months; the placebo group wore similar, unmagnetized device.

**Intervention:** Nerve conduction and/or quantified sensory testing were performed serially.

**Main Outcome Measures:** Daily visual analog scale scores for numbness or tingling and burning and QOL issues were tabulated over 4 months. Secondary measures included nerve conduction changes, role of placebo, and safety issues. Analysis of variance (ANOVA), analysis of covariance (ANCOVA), and chi-square analysis were performed.

**Results:** There were statistically significant reductions during the third and fourth months in burning (mean change for magnet treatment,  $-12\%$ ; for sham,  $-3\%$ ;  $P < .05$ , ANCOVA), numbness and tingling (magnet,  $-10\%$ ; sham,  $+1\%$ ;  $P < .05$ , ANCOVA), and exercise-induced foot pain (magnet,  $-12\%$ ; sham,  $-4\%$ ;  $P < .05$ , ANCOVA). For a subset of patients with baseline severe pain, statistically significant reductions occurred from baseline through the fourth month in numbness and tingling (magnet,  $-32\%$ ; sham,  $-14\%$ ;  $P < .01$ , ANOVA)

and foot pain (magnet,  $-41\%$ ; sham,  $-21\%$ ;  $P < .01$ , ANOVA).

**Conclusions:** Static magnetic fields can penetrate up to 20mm and appear to target the ectopic firing nociceptors in the epidermis and dermis. Analgesic benefits were achieved over time.

**Key Words:** Diabetic neuropathies; Magnetics; Rehabilitation.

© 2003 by the American Congress of Rehabilitation Medicine and the American Academy of Physical Medicine and Rehabilitation

**D**IABETIC PERIPHERAL NEUROPATHY (DPN) is a common and often disabling complication of diabetes mellitus (DM). Depending on criteria, DPN is estimated to occur in 50% to 90% of individuals with diabetes for more than 10 years.<sup>1-4</sup> As many as half of the 16 million diabetics in the United States will experience neuropathic pain at some point in their lives.<sup>5-9</sup> DPN begins insidiously, presenting as a symmetrical sensory polyneuropathy that follows a stocking-glove pattern. Selective involvement of unmyelinated C fibers and small myelinated A delta fibers produces pain of the burning dysesthetic type and is often accompanied by hyperalgesia and allodynia in the feet.<sup>7,10-12</sup> Neuropathic pain symptoms fluctuate and can be described as superficial, deep, aching, lancinating, constant, or episodic. Complaints are often worse at night. Although initial symptoms and the course of DPN vary, once neuropathic pain is established, it is almost always progressive, leading to increased discomfort and disability.<sup>6,13-15</sup> Furthermore, individuals with DPN are at augmented risk for foot trauma and infections that may necessitate amputative procedures.<sup>2,16</sup>

From a pathophysiologic standpoint, these symptoms are believed to be secondary to ectopic firing of nociceptive afferent axons that are undergoing degeneration.<sup>7,9-12</sup> This ectopic depolarization appears to be related to dysregulated expression of sodium and calcium channels<sup>17-19</sup> and a deficit in the potassium-internal rectifying channel.<sup>20-22</sup> Neurons at the level of the dorsal root ganglion (DRG) also become hyperexcitable after peripheral nerve injury, presumably because of loss of peripheral inhibitory influences.<sup>23</sup> Currently, there are no treatments that reverse or arrest progressive diabetic polyneuropathy.<sup>24</sup> A variety of standard oral therapies used for symptomatic neuropathic pain include tricyclic antidepressants,<sup>25</sup> antiepileptic medications,<sup>26</sup> and narcotic analgesics.<sup>27,28</sup> Additionally, topical products such as capsaicin<sup>29,30</sup> have been applied and have produced incomplete pain relief and significant side effects. Overall, the results have been disappointing and associated with significant side effects.<sup>15,31,32</sup> The search for reliable, safe, and effective mainstream treatments for the neuropathic pain of DPN remains a major challenge,<sup>13,15,25-27,31-34</sup> and, not surprisingly, patients have explored a variety of alternative approaches, including homeopathy, acupuncture, and magnetic

From the Department of Neurology, New York Medical College, Valhalla, NY (Weintraub); University of Texas, Southwestern Medical Center, Dallas, TX (Wolfe, Barohn); Research Design Inc, Yorktown Heights, NY (Cole); University of Minnesota, Minneapolis, MN (Parry); St. Louis University, St. Louis, MO (Hayat); Kaiser-Permanente Medical Group, Denver, CO (Cohen); California College of Podiatric Medicine, San Francisco, CA (Page); University of Utah, Salt Lake City, UT (Bromberg); and Diabetes and Glandular Disease Clinic, San Antonio, TX (Schwartz).

Supported by Nu-Magnetics Inc and Nikken Inc.

Presented in part at the Joint Conference of the American Congress of Rehabilitation Medicine and the American Society of Neurorehabilitation, October 5, 2002, Philadelphia, PA.

No commercial party having a direct financial interest in the results of the research supporting this article has or will confer a benefit upon the author(s) or upon any organization with which the author(s) is/are associated.

Reprint requests to Michael I. Weintraub, MD, Dept of Neurology and Medicine, New York Medical College, 325 S Highland Ave, Briarcliff Manor, NY 10510, e-mail: [miwneuro@pol.net](mailto:miwneuro@pol.net).

0003-9993/03/8405-736\$30.00/0

doi:10.1016/S0003-9993(03)00106-0

therapies. Spurred on by anecdotal reports, the use of permanent magnets for relief of pain has become extremely popular in recent years, with consumer spending exceeding \$500 million in the United States and Canada and \$5 billion worldwide.<sup>35,36</sup> The idea that magnetic energy from commercially available, weak magnets applied locally to the feet could influence chronic neuropathic pain may seem absurd, and yet this approach is not new.<sup>37-41</sup> In the absence of randomized, placebo-controlled trials, the medical community has been understandably skeptical, which has limited the acceptance of magnets as a valid option for pain relief.<sup>42,43</sup> However, 2 prior pilot studies successfully showed reduced neuropathic pain in 75% and 90% of patients with refractory DPN over a 4-month period, with constant application of commercial multipolar foot magnets (450G).<sup>35,36</sup> These surprising and unexpected favorable results prompted the present study—a nationwide, randomized placebo-controlled investigation into the legitimacy of static magnetic fields in the relief of pain from DPN.

## METHODS

### Enrollment Criteria

From August 1999 through January 2001, 375 subjects with symptomatic symmetrical sensory and motor diabetic peripheral neuropathy (DPN stages II or III), as defined by Dyck et al,<sup>44,45</sup> were recruited from 48 sites in 27 states. Consecutive patients from neurologic, podiatric, and diabetic clinics or private practice were enrolled. A few centers advertised their participation in this nationwide study to attract eligible volunteers. The primary providers were skilled clinicians who had previously participated in pharmacologic studies of diabetes and/or pain management. Enrollment criteria required that all subjects have at least 2 abnormalities on neurologic examination (sensory, motor, reflex), moderate (II) to severe (III) neuropathic pain, abnormal nerve conduction or quantitative sensory testing, and/or symptoms of autonomic dysfunction. Symptoms had to be constant and present over 6 months and refractory to various medications. Subjects included persons with insulin-dependent diabetes mellitus (IDDM) and those who were not insulin dependent (NIDDM). Subjects were excluded if other systemic diseases could potentially explain their symptoms. As a safety precaution, pregnant women and subjects who had mechanical insulin pumps or cardiac pacemakers were also excluded. Subjects tabulated validated<sup>46-50</sup> daily pain scores and similar, but unvalidated, quality of life (QOL) scores for 4 months and agreed that they would not attempt to break blinding of the foot devices. They also agreed to wear the devices constantly, 24 hours per day. Moderate pain was defined as scores of 5.0 to 6.99 and severe pain was defined as 7 and higher. No new analgesic drugs were allowed during the study, but individuals could remain on (or reduce) their current regimen of neuropathic pain medication. The randomized, placebo-controlled, parallel design study was fully explained to all subjects and voluntary withdrawal was allowed without prejudice.

### Randomization

Demographic data (age, height, weight, gender, race, glycosylated hemoglobin [Hb A<sub>1c</sub>], family history, duration of DM, complications of DM, treatment of DM) were collected at each site. Subjects completed a 2-week baseline Likert visual analog scale (VAS) quantification of their pain symptoms 3 times daily to establish a reliable mean pain score. QOL scores were recorded once daily to measure (1) sleep disturbance secondary to foot pain and (2) exercise-induced foot pain after a 10-

minute exertion such as walking or other physical activity. After eligibility was confirmed and written informed consent accepted, subjects were randomized consecutively (1:1 via computer assignment) to receive an active magnetic shoe insole or a sham insole of similar appearance. Randomization was stratified by center and gender. Neither the subject nor the research staff was aware of the treatment allocation. If corrective trimming of the device was necessary to provide a comfortable fit in the shoe, a noninvolved secretary or nurse would trim them along identifiable lines around the margins. The subjects and site investigators were not present if trimming was necessary. All data were submitted to a central data bank under the supervision of the statistician who was aware of the assignments.


### Magnetic Devices

The devices used in the present study are comprised of a reinforced and flexible magnetic rubber compound pressed into a sheet and cut into the shape of a shoe insole for men and women. Strontium ferrite powder is mixed into this rubber binder and magnetized with a patented pattern of alternating magnetic poles. Each pole is adjacent to and contiguous with another triangular-shaped magnetic pole of opposite polarity on each of the 3 sides of the triangle. This pattern produces a continuous array of alternating magnetic poles in every direction across the insole (fig 1).

The strength of the magnetic field is 450G, as measured with a conventional gauss meter on the surface of the insoles at the center of the triangle (10,000G=1T). The field depth of penetration is 20mm and is reduced inversely with the square of the distance. By far, the simple, most direct method of determining field strength at various distances from the insole surface is by instrument measurement. For example, using a Lakeshore 420 gauss meter with a flat transverse probe<sup>a</sup> has an accuracy of  $\pm .25\%$ . The effective field of the magnet from the insole surface is 20mm. Beyond 20mm, the magnetic field measures in the range of the ambient magnetic field of the earth at about 0.5G. The maximum surface field strength of the magnetic insole is 450G. At a 1-mm distance from the surface, the field strength drops to 249G. At 2mm, the field strength is measured at 150G. At 3mm (approximately  $\frac{1}{8}$ in), the field strength is 90G. Flux density at the target area may be more clinically relevant than the magnetic reading at the surface of the magnet. The specific flux density, however, at the target area is unknown. At 13mm above the surface of the magnetized insole, the reading is only 1.5G. The sham insole's gauss meter readings did not exceed the 0.5G of the earth's magnetic field. Both sham and active magnetic shoe insoles could not be distinguished in terms of appearance, consistency, or weight. The magnetic insoles used in the present study were manufactured by Nu-Magnetics Inc,<sup>b</sup> and are commercially sold under the brand name of Magsteps<sup>®</sup> by Nikken Inc.<sup>c</sup>

### Outcome Measures

Pain was measured on an 11-point numeric pain rating scale (VAS; scale range: 0, no pain; 10, worse possible pain). The primary efficacy measure was the reduction in neuropathic pain scores at week 16 compared with baseline scores. We also compared month-to-month changes. We looked specifically at 2 of the most common pain symptom scores of numbness or tingling and burning. Each symptom was recorded 3 times daily so to reduce any new variables (VAS range, 0-10). Similarly, QOL issues were considered primary efficacy measures with reduction of exercise-induced foot pain and sleep interruption secondary to pain (VAS range, 0-10). These were recorded once daily. Secondary outcomes compared baseline



**Fig 1.** Magnetic field visualization with superimposed magne-view film. The microencapsulated colloidal nickel particles congregate in alignment with the magnetic flux lines producing a 2-dimensional image of the pole pattern.

and 16-week values of neurologic examinations, nerve conduction velocity (NCV), quantitative sensory testing (QST) thresholds (Neurometer<sup>®51d</sup> or Case IV<sup>52</sup>), and other electrophysiologic tests.<sup>53,54</sup> Safety measures with tabulation of adverse events were monitored as was cause for dropouts. Additionally, an interim study performed before the end of this study at selected sites assessed masking and bias by asking patients and investigators whether they believed that a placebo or active device was used or whether they had no opinion.

### Sites

There were 48 investigative sites in 27 states. They included 11 university-based centers and 37 private practices. A neurologic examination was performed before entry to identify the presence of a sensory peripheral polyneuropathy in the feet that met the Dyck<sup>45</sup> criteria of moderate (II) to severe (III) DPN. NCVs of the peroneal and/or posterior tibial (motor) and sural nerves (sensory) were performed in a standardized manner to confirm the presence of neuropathy. Selected sites performed forced-choice QST by using Neurometer (CPT) or Case IV equipment and other neurophysiologic tests, such as biothesiometry and sympathetic skin response (SSR). Because no standard, validated device exists and controversy about their merits surrounds the various devices, we let each site use their standard analysis technique.

### Investigational Review Board

Phelps Memorial Hospital Investigational Review Board (IRB) reviewed and approved the protocol, as did IRBs at individual university centers. Phelps Memorial served as a central IRB for many investigative sites and appropriate safety and progress data were submitted to this IRB in a timely fashion. All patients provided written informed consent to participate in this study.

### Statistical Analyses

For each of the 4 outcome measures (burning, numbness and tingling, foot pain, sleep scores), a 2 (treatment, sham)  $\times$  5 (baseline, 1mo, 2mo, 3mo, 4mo) repeated-measures analysis of variance (ANOVA) was used to assess possible differences between treatment and sham groups over the course of the study. These analyses were followed by a 2 (treatment, sham)  $\times$  2 (2mo, 4mo) analysis of covariance (ANCOVA) with baseline score as the covariate to explore treatment effects during the last 2 months of the study. Furthermore, for each outcome measure, we grouped patients into 3 categories of severity based on baseline scores. Ratings of 1 to 4 corresponded to mild pain; 5.0 to 6.99, to moderate pain; and 7 to 10, to severe pain.<sup>55</sup> ANOVAs were used to compare the mean changes separately for each severity group. For each of the outcome measures, chi-square tests for independence were used to assess magnet versus sham group differences in the percentage of patients who had at least a 30% reduction in severe pain. Finally, ANOVAs and ANCOVAs were used to assess treatment effects for subgroups defined by measures known to previously affect outcomes in this population. For all tests, a *P* value of .05 or less was considered to indicate statistical significance. Subjects with any missing data for an endpoint were excluded for that analysis.

On the basis of published results of clinical trial placebo responses for painful diabetic neuropathy,<sup>26</sup> at an  $\alpha$  level of .05 and a power of .80, with 150 subjects per group, it was estimated that a difference between treatment and sham group responses of 17% or more would be statistically significant.<sup>56</sup> Analyses were conducted with SPSS.<sup>e</sup>




Fig 2. Flowchart of the randomized placebo-control trial. Abbreviation: Rx, treatment.

### Adverse Events

Potential injury to the sole producing ulcer or abrasion or infection was monitored. Mechanical allodynia because of sensitive feet was also tabulated.

### Role of Funding Source

This study was initially funded by Nu-Magnetics and supplemented by Nikken Inc. The grant recipients had complete independence regarding study design, data analysis, and manuscript preparation. The study's protocol was approved by the National Institutes of Health, but not funded.

## RESULTS

The flow of patients through the clinical trial is depicted in figure 2. Three hundred seventy-five subjects were randomly assigned to treatment and sham groups, and 259 subjects (69%) successfully completed this 4-month trial. Of the 90 dropouts, 74% in the treatment group and 71% in the sham cohort dropped out before the second month. Of the total group, 45% were lost to follow-up, 24% dropped because of allodynia, and 9% dropped for nonstudy complications. Twenty-six subjects were dropped by the statistician for missing or questionable

data. The baseline characteristics for the remaining 259 subjects were similar for treatment and sham groups (table 1). The *t* tests for independent samples revealed no baseline differences between the treatment and sham groups for the primary end points (table 2). Racial-ethnic proportions at enrollment were a representative cross-section of the US population. In addition, a series of ANOVAs revealed no baseline differences or differences over the study period between patients at university centers and in private practice settings.

### Primary Outcomes

**Burning.** Burning scores decreased 30% for the treatment group from baseline (mean  $\pm$  standard deviation,  $5.13 \pm 2.29$ ) to month 4 ( $3.61 \pm 2.44$ ) and decreased 24% for the sham group from baseline ( $5.27 \pm 2.40$ ) to month 4 ( $4.01 \pm 2.81$ ) ( $P = .000$ , ANOVA; fig 3). There was a larger decrease in mean scores for the treatment group ( $-12\%$ ) from month 2 ( $4.09 \pm 2.38$ ) to month 4 ( $3.61 \pm 2.44$ ) than for the sham group ( $-3\%$ ) from month 2 ( $4.12 \pm 2.65$ ) to month 4 ( $4.01 \pm 2.81$ ) ( $P < .05$ , ANCOVA).

**Numbness and tingling.** Numbness and tingling scores decreased 29% for the treatment group from baseline

Table 1: Baseline Characteristics of the Subjects

| Characteristic                | Treatment Group (n=141) | Sham Group (n=118) |
|-------------------------------|-------------------------|--------------------|
| Age (y)                       |                         |                    |
| Mean                          | 62.6±11.3               | 63.2±11.2          |
| Range                         | 36-85                   | 27-85              |
| Weight (lb)                   | 206.7±47.0              | 207.1±41.2         |
| Height (in)                   | 67.7±4.05               | 67.9±4.28          |
| Sex (n)                       |                         |                    |
| Female                        | 66                      | 58                 |
| Male                          | 75                      | 60                 |
| Race (n)                      |                         |                    |
| White                         | 107                     | 103                |
| Nonwhite                      | 34                      | 15                 |
| Years since onset of diabetes | 13.0±10.8               | 11.6±10.2          |
| HB A <sub>1c</sub>            | 7.7±1.8                 | 7.6±2.1            |
| Nerve conduction velocity (n) |                         |                    |
| Normal                        | 5                       | 3                  |
| Axonal                        | 42                      | 31                 |
| Demy                          | 16                      | 14                 |
| Mixed                         | 51                      | 49                 |
| Insulin (n)                   |                         |                    |
| Yes                           | 49                      | 40                 |
| No                            | 92                      | 78                 |

NOTE. Values are mean ± standard deviation (SD) or as otherwise indicated. Abbreviation: Demy, demyelinating.

(5.63±2.08) to month 4 (4.02±2.46) and decreased 22% for the sham group from baseline (5.89±2.02) to month 4 (4.57±2.58) ( $P=.000$ , ANOVA; fig 4). There was a decrease in mean scores for the treatment group (-10%) from month 2 (4.46±2.23) to month 4 (4.02±2.46) and a small increase for the sham group (+1%) from month 2 (4.54±2.58) to month 4 (4.57±2.58) ( $P<.05$ , ANCOVA). For patients with severe pain at baseline, numbness and tingling decreased 32% for the treatment group from baseline (8.17±.85) to month 4 (5.58±2.43) and decreased 14% for the sham group from baseline (8.12±.95) to month 4 (6.97±2.38) ( $P<.01$ , ANOVA; fig 5). Of the 38 treatment patients with severe pain at baseline, 27 (71%) had mild or moderate pain at month 4. In contrast, of the 40 sham patients with severe pain at baseline, 16 (40%) had mild or moderate pain at month 4 ( $P<.01$ ,  $\chi^2$ ).




Fig 3. Burning mean scores for treatment and sham subjects.

**Foot pain.** Foot pain scores decreased 31% for the treatment group from baseline (5.84±2.33) to month 4 (4.05±2.66) and decreased 25% for the sham group from baseline (5.76±2.29) to month 4 (4.31±2.80) ( $P=.000$ , ANOVA; fig 6). A larger decrease in mean scores existed for the treatment group (-12%) from month 2 (4.62±2.53) to month 4 (4.05±2.66) than for the sham group (-4%) from month 2 (4.47±2.68) to month 4 (4.31±2.80) ( $P<.05$ , ANCOVA). For patients with severe pain at baseline, foot pain decreased 41% for the treatment group from baseline (8.49±1.07) to month 4 (4.97±3.10) and decreased 21% for the sham group from baseline (8.35±.95) to month 4 (6.56±2.50) ( $P<.01$ , ANOVA; fig 7). Of the 40 treatment patients with severe pain at baseline, 29 (69%) had mild or moderate pain at month 4. In contrast, of the 35 sham-device patients with severe pain at baseline, 17 (49%) had mild or moderate pain at month 4. This trend in category change did not reach statistical significance ( $P=.07$ ,  $\chi^2$ ).

**Sleep.** Sleep scores decreased 30% for the treatment group from baseline (4.83±2.66) to month 4 (3.36±2.76) and decreased 30% for the sham group from baseline (5.19±2.79) to month 4 (3.65±3.04) ( $P=.000$ , ANOVA; fig 8). There was a nonsignificant trend for a larger decrease in mean scores for the treatment group (-13%) from month 2 (3.83±2.83) to month

Table 2: Mean Scores for Primary Endpoints From Baseline to Month 4

| Outcome Measure              | n   | Baseline | Month 1 | Month 2 | Month 3 | Month 4 |
|------------------------------|-----|----------|---------|---------|---------|---------|
| <b>Burning</b>               |     |          |         |         |         |         |
| Treatment                    | 133 | 5.1±2.3  | 4.3±2.3 | 4.1±2.4 | 3.9±2.5 | 3.6±2.4 |
| Sham                         | 111 | 5.3±2.4  | 4.6±2.6 | 4.1±2.7 | 4.1±2.7 | 4.0±2.8 |
| <b>Numbness and tingling</b> |     |          |         |         |         |         |
| Treatment                    | 137 | 5.6±2.1  | 4.7±2.2 | 4.5±2.2 | 4.3±2.4 | 4.0±2.5 |
| Sham                         | 116 | 5.9±2.0  | 4.9±2.3 | 4.5±2.6 | 4.6±2.6 | 4.6±2.7 |
| <b>Foot pain</b>             |     |          |         |         |         |         |
| Treatment                    | 121 | 5.8±2.3  | 4.9±2.4 | 4.6±2.5 | 4.2±2.6 | 4.1±2.7 |
| Sham                         | 106 | 5.8±2.3  | 4.9±2.4 | 4.5±2.7 | 4.3±2.8 | 4.3±2.8 |
| <b>Sleep</b>                 |     |          |         |         |         |         |
| Treatment                    | 112 | 4.8±2.7  | 4.0±2.8 | 3.8±2.8 | 3.5±2.7 | 3.4±2.8 |
| Sham                         | 98  | 5.2±2.8  | 4.6±2.6 | 3.8±2.8 | 3.8±3.0 | 3.7±3.0 |

NOTE. Values are mean ± SD.




Fig 4. Numbness and tingling mean scores for treatment and sham subjects.

4 ( $3.36 \pm 2.76$ ) than for the sham group ( $-3\%$ ) from month 2 ( $3.76 \pm 2.83$ ) to month 4 ( $3.65 \pm 3.04$ ) ( $P = .08$ , ANCOVA).

**Secondary Outcomes**

There was no evidence of deterioration of nerve function clinically or electrophysiologically in those patients reporting improvement in pain scores. Thus, there was no evidence of clinical worsening. Of the 259 subjects, 61 (24%) had Neurometer, Case IV, SSR, or biothesiometry studies. No significant differences existed between subjects in the treatment group ( $n=32$ ) and those in the sham group ( $n=29$ ) from baseline to 4 months on these measures.

**Subgroup Analyses**

For patients not taking oral antidiabetic agents, a larger decrease occurred in mean burning scores for the treatment group ( $-14\%$ ) from month 2 ( $3.81 \pm 2.38$ ) to month 4 ( $3.30 \pm 2.39$ ) than for the sham group ( $-1\%$ ) from month 2 ( $3.91 \pm 2.87$ ) to month 4 ( $3.86 \pm 2.85$ ) ( $P < .01$ , ANCOVA). There was a nonsignificant trend for a larger decrease in mean numbness and tingling scores for the treatment group ( $-10\%$ )




Fig 5. Numbness and tingling mean scores for subjects with baseline severe pain.




Fig 6. Foot pain mean scores for treatment and sham subjects.

from month 2 ( $4.26 \pm 2.21$ ) to month 4 ( $3.84 \pm 2.46$ ) than for the sham group ( $-1\%$ ) from month 2 ( $4.78 \pm 2.68$ ) to month 4 ( $4.24 \pm 2.59$ ) ( $P = .08$ , ANCOVA). A similar pattern was reported for patients with severe foot pain scores, with reductions of 41% and 21% for treatment and sham groups, respectively, and for numbness and tingling, with reductions of 32% and 23% for the 2 groups, respectively. Results remained significant with a Bonferroni correction.<sup>57</sup> By using the 30% pain reduction criterion as suggested by a Farrar stratification analysis,<sup>58</sup> we noted that 50% of patients with magnets had at least a 30% reduction in severe numbness and tingling, compared with 25% of patients with sham devices ( $P < .05$ ,  $\chi^2$ ). Although the percentages for foot pain (32% vs 19%) and burning (42% vs 29%) were impressive, they were not statistically significant. No differences between treatment and sham groups were found based on family history of diabetes, baseline nerve conduction, or Hb A<sub>1c</sub> scores.

**Blinding**

An interim analysis for bias and breaking the blind was performed at those active sites 6 months before study terminated (university and private practice). This analysis was to determine whether the present study was adequately blinded.




Fig 7. Foot pain mean scores for subjects with baseline severe pain.




Fig 8. Sleep mean scores for treatment and sham subjects.

Subjects and examining investigators were asked at the end of the study to identify the treatment provided. Sixty-three percent of the subjects responded. Of the 83 treatment group subjects responding, 40 (48%) believed they had active magnets, 31 (37%) believed they had sham magnets, and 12 (15%) did not know. Of the 80 sham-device subjects responding, 29 (36%) believed they had active magnets, 30 (38%) believed they had sham magnets, and 21 (26%) did not know. Of 46 investigators of treatment subjects, 23 (50%) believed the subjects had active magnets, 15 (33%) believed they had sham magnets, and 8 (17%) did not know. Of 50 investigators of sham-device subjects, 22 (40%) believed the subjects had active magnets, 15 (30%) believed they had sham magnets, and 12 (26%) did not know. There was no significant association between the actual treatment received and the belief about the treatment received for subjects or investigators.

### Dropouts

The dropouts were evenly represented and did not impact on the primary analysis for efficacy. We did not use the intention-to-treat (ITT) model for estimates of missing data, because 75% of the dropouts from the treatment group and 71% from the sham group dropped out before month 2. As shown in our figures, the magnetic effects became apparent after month 2; therefore, using the ITT model with most estimates based on data before month 2 would severely bias the analysis. Dropouts secondary to allodynia were equally common in both groups. Foot sensitivity is a well-known phenomenon in symptomatic patients with DPN. Thus, it is not surprising that the application of an insole (magnetized or unmagnetized) would be unpleasant to a small but significant group of patients. There were 90 dropouts (lost to follow-up, allodynia, complications) equally represented out of a sample size of 349 (25.8%). There were no mean differences between the 46 treatment and 44 sham-device patients for age, years since onset of diabetes, and baseline Hb A<sub>1c</sub>, burning, numbness and tingling, foot pain, and sleep scores ( $P > .05$ , ANOVA). The statistician dropped 26 patients (equal representation) because of site difficulties obtaining data and unreliable data.

### Safety

Measures of safety included constant reporting of adverse events and the cause for dropouts. There were no significant complications.

## DISCUSSION

This is the first multicenter, double-blind, placebo-controlled study to examine the role of static magnetic fields in a homogeneous cohort of DPN with neuropathic pain. The antinociceptive effect was significantly pronounced during the third and fourth month, indicating that a tonic and chronic exposure must be present to inhibit and influence sensitized afferent pain fibers. The magnitude of the reduction of burning, numbness and tingling, and exercise-induced foot pain, especially in severe and extreme cases, was comparable or superior to that observed in the gabapentin,<sup>26</sup> tramadol,<sup>28</sup> and lamotrigine<sup>24</sup> studies, but without side effects. Additionally, a change of 1.5 in the 0 to 10 pain scale represents a clinically meaningful difference.<sup>59,60</sup> This also reaffirms the data from 2 prior pilot studies.<sup>35,36</sup> Subset analysis identified that subjects with severe pain<sup>55</sup> and those not taking oral hypoglycemic agents responded more favorably than other symptomatic patients. Although our results show a statistically significant reduction in predetermined primary outcome measures, it is difficult to determine the mechanism of action responsible for these benefits. It is of interest that in the pharmacologic trials of tramadol<sup>28</sup> and gabapentin,<sup>26</sup> the subjects with severe and extreme pain responded better than other subjects. Segal et al<sup>61</sup> also noted in testing bipolar magnetic devices in knee pain secondary to rheumatoid arthritis that patients with mild symptoms did not respond as well. DPN pain appears to arise from an increase in afferent signals from degenerating nociceptive afferent fibers. It has been shown that early in the course of painful neuropathies, free nerve endings of nociceptive axons can disappear from the skin but are still present in the sural nerve.<sup>62</sup> One possibility may be that the magnetic field of these insoles somehow directly or indirectly interrupts and suppresses the afferent signal traffic of the C-fiber firing pattern of the distal part of the surviving axon thereby producing an antinociceptive effect. A number of studies have shown that DPN pain could result from depolarization because of dysregulation of normal sodium,<sup>17-19,63</sup> calcium,<sup>23,64</sup> and potassium<sup>20</sup> channel activities. It is well known that sodium channels accumulate in areas of axonal damage<sup>63</sup> and static magnetic fields have been shown to block or reduce action potential via effects on sodium flux.<sup>65-68</sup> A number of studies using weak pulsed, time-varying electromagnetic fields have shown biologic changes.<sup>69-73</sup> Adey and Chopart<sup>74,75</sup> considered the cell membrane as the most likely transducer modifying ion transport of protein and adenosinetriphosphatase activity. Membrane lipids with organized arrays of polar molecules, diamagnetic, have been shown to realign anisotropic molecules as well as to summate and interfere with ionic transport.<sup>76,77</sup> Translational movement or changes in orientation in a magnetic field can influence amplitude of evoked responses.<sup>78,79</sup> Because phospholipids in cell membranes have both diamagnetic and paramagnetic properties, it is clear that mechanisms exist that can produce conformational changes in various channels and structures.<sup>80,81</sup> However, it is not known if any of this is pertinent to putative biologic effects of static magnetic fields. Based on our data, we speculate that the kinetic activity of channelized membrane ions and blood flow in a static magnetic field is sufficiently strong to stimulate living tissues and to induce a biologic reaction. Signal transduction pathways appeared to be functionally modulated, and this is a restatement of Faraday's law of time variation.<sup>70,82,83</sup> It is also known that weak magnetic fields can increase the partial pressure of tissue oxygen, thereby improving oxygen delivery to tissues.<sup>84</sup> This property may be important because of a reported reduction in endoneurial oxygen tension in DPN.<sup>85</sup> Thus, it is biologically plausible

that static magnetic fields influence diabetic neurons and cell membranes of cutaneous nociceptors by amplifying the weak electromagnetic signals from the imposed and constant static magnets, thereby inducing changes in the cellular<sup>86-88</sup> and pericellular microenvironment.<sup>89,90</sup> Because these devices have a presumed penetration of up to 20mm—thereby indicating passage through the epidermal<sup>91</sup> and dermal layers, which contain a rich network of nerves and capillaries—we speculate that, at this site, there is inhibition and/or interruption of ectopic firing of the damaged small nociceptive afferent unmyelinated C fibers. The specific magnetic flux density at this target area is not known. Perhaps a gating response with simultaneous stimulation of the A delta fibers producing an inhibitory antinociceptive effect on C fibers occurs, compatible with Melzak–Wall hypothesis.<sup>92</sup> Another possibility includes the recruitment of previously passive C fibers.<sup>93,94</sup> Case IV studies of warm and/or cold thermal thresholds did not reveal any serial changes from baseline. Thus, at an ionic-membrane level, we can speculate that either the underlying sodium channels can be up- or down-regulated<sup>95</sup> or, alternatively, rapid repolarization occurs because of stimulation of the potassium internal rectifying channels.<sup>64</sup> This phenomenon may also produce a secondary inhibition of the firing from the DRG neurons.<sup>23</sup>

The major strengths of the present study include randomized, placebo design; the cooperative involvement of neurologists, podiatrists, and diabetologists; and the geographic and racial diversity of the study population. These factors suggest that the observed benefits will be applicable to the general diabetic population. Because pain levels can vary during the day, patients recorded their score 3 times daily to best derive a mean daily discomfort level and to reduce recall bias. Similarly, QOL experiences have yet to be standardized and validated by large cohorts in DPN<sup>34</sup>; yet, intuitively, quantification of exercise-induced foot pain and sleep disturbance represents important functional outcome measures.<sup>96,97</sup> Another strength is the utilization of both academic and private practice centers that not only showed good interobserver reliability, but also reduced the likelihood of selection bias.

Despite this provocative data, several limitations exist. We relied exclusively on patients' self-report for pain and outcome.<sup>55,98</sup> Despite favorable statistical reduction of neuropathic pain and QOL scores by wearing these devices, only modest clinical improvement was achieved. The slopes of our figures from months 2 to 4 suggest that a more potent clinical benefit could be anticipated at 8 to 12 months, and, thus, long-term studies must be performed. Another limitation was that it is a physical impossibility to blind these foot devices and to prevent the determination of magnetic activity. Subjects and investigators were advised of the importance of maintaining the blind, and the questionnaire at study termination indicates that both groups remained blinded.<sup>99,100</sup> Unfortunately, we were unable to identify a biologic marker using QST, SSR, and biothesiometry. None of the limitations invalidates the statistical antinociceptive effects. Intraepidermal nerve fiber density measurements were not performed and may have provided a useful pathologic correlate.<sup>101</sup> It has been shown that regeneration of nerve fibers can occur within 39 days in the dermis after an injury and after 4 months in the epidermis.<sup>102,103</sup> The observation that both refractory groups improved with lower VAS scores by 2 months compared with baseline by wearing foot devices (magnetized, unmagnetized) is provocative and similar to that seen in pharmaceutical studies and placebo trials; this suggests either a placebo response or analgesic benefit induced by foot pressure. It is possible that central regions of the brain

for pain control (ie, rostral anterior cingulate cortex, brainstem) were somehow activated.<sup>29</sup>

## CONCLUSION

Although many questions remain about a precise mechanism of action, the present study provides convincing data confirming that the constant wearing of static, permanent, magnetic insoles produces statistically significant reduction of neuropathic pain. Considering their safety and minimal cost (<\$100), our data suggest that the insoles may be used as adjunctive or monotherapy. Future studies are needed to identify the optimal time to achieve maximum antinociceptive effect and to confirm and extend these results. Additional search for biologic markers (ie, epidermal nerve fiber biopsy, microneurography) will be necessary in future protocols to determine if permanent structural changes can be produced.

**Acknowledgments:** We thank Jeffrey Katims, MD, of Neurotron Inc, for the loan of equipment used at the 3 centers. We also thank Dr. William Frishman for critical review of the manuscript and Susan Pines-Wolert for data collection.

The members of the Magnetic Research–Diabetic Neuropathy Study Group include the following: M.I. Weintraub, MD (principal investigator); A.L. Rothman, MD, and G.I. Wolfe, MD (co-principal investigators); and Steven P. Cole, PhD (chief biostatistician).

Site investigators (listed alphabetically with the principal investigator listed first) include the following: B.T. Adornato, MD, P.C. Cassini, MD, Stanford Medical Center, Palo Alto, CA; C.N. Applegate, MD, E. Moore, CNRN, Ozarks Medical Center, West Plains, MO; S.W. Asher, MD, Neurological Associates, Boise, ID; T.E. Bertorini, MD, J. Karb, LPN, S. Maccarino, R.EDT, Wesley Neurology Clinic, Memphis, TN; T.W. Bohr, MD, V. Johnson, MD, D. Moses, RN, Loma Linda University School of Medicine, Loma Linda, CA; M.B. Bromberg, MD, A.K. Faucher, RN, University of Utah Medical Center, Salt Lake City, UT; H.K. Bucholtz, MD, R. Malone, RN, Edison, NJ; A.C. Chalmers, MD, S. Tipton, Kern County Neurological Medical Group, Bakersfields, CA; R. Cintron, Neuroscience Consultants, Reston, VA; J.A. Cohen, MD, F. Zeren, RN, MSN, D. Armbruster, RN, Kaiser Permanente Medical Group, Denver, CO; S.P. Cole, PhD, Research Design Associates, Yorktown Heights, NY; S. Cooper, MD, Chad Breaux, CRC, Medical Associates of Georgia, Canton, GA; A.C. Cuetter, MD, O. Molinar, RN, Texas Tech Medical Center, El Paso, TX; N.W. Culligan, MD, S. Lindblom, PT, Associated Neurologists PC, Danbury, CT; L. Diamond, MD, New York College of Podiatric Medicine, New York, NY; P.B. Dunne, MD, S. Khoromi, MD, H. Wang, MD, University of South Florida, Tampa, FL; L.W. Epperson, MD, S. Thorp, RN, Alabama Neurological Clinic, Montgomery, AL; A.J. Esposito, MD, Northeast Alabama Neurological Services, Anniston, AL; R.A. Fischer, MD, L.M. Leschek-Gelman, MD, R. Titus, RN, Neurology Associates, Wilmington, DE; D.F. Fleming, MD, M.T. Stock, DPhil, FNP, East Carolina Neurology, Greenville, NC; T. Giancarlo, DO, G. Lapadot, LPN, Michigan Neurology Associates, St. Claire Shores, MI; M.G. Gregory, MD, J. Bishop, RN, CCRC, Nevada Neurological Consultants, Las Vegas, NV; I. Haber, DO, M. Smith, Terre Haute, IN; G. Hayat, MD, J. Armbruster, RN, St. Louis University, St. Louis, MO; S.A. Kabbani, MD, T. Jenkins, East Tennessee Neurology Clinic, Knoxville, TN; J. Kawalec, PhD, B.D. Caldwell, DPM, A. Patel, MD, Ohio College of Podiatric Medicine, Cleveland, OH; H.L. Kettler, MD, FAAN, Wheeling Clinic, Wheeling, WV; D.A. Konanc, MD, K.L. Hull, MD, A.T. Perkins, MD, S.M. Freedman, MD, P.F. Bye, RN, CRC, J.K. Downs, RN, CCRC, T.S. Garriss, CRC, Raleigh Neurology Associates, Raleigh, NC; K.A. Levin, MD, K.A. Citak, MD, J.T. Nasr, MD, S. Pipala, RN, L. Warnock, RN, Neurology Group of Bergen County, Ridgewood, NJ; M.E. Lipitz, DO, L.L. Ford, LPN, J.E. Benton, RN, Blair Medical Associates, Altoona, PA; A. Maloon, MD, D. Hall, LPN, CCRC, Marietta Neurological Associates, Marietta, GA; R. Mendicino, DPM, D. Haupt, RN, Foot and Ankle Institute of Western Pennsylvania, Pittsburgh, PA; C. Miller, MD, B.C. Wouters, MD, D.D. Mayer, RN, Neurology Consultants of Montgomery, Montgomery, AL; J. Page, DPM, CH DPM, California College of Podiatric

Medicine, San Francisco, CA; G.L. Pattee, MD, D. Hartmann, RN, Neurology Associates, Lincoln, NE; T.A. Payne, MD, J. Romanowsky, MD, R. Antil, LPN, Neurology Clinic of Saint Cloud, Saint Cloud, MN; T.J. Regan, MD, C.B. Ward, CCR, Hampton Roads Neurology, Newport News, VA; S. Saeed, MD, V. Prater, LPN, West Tennessee Neurology, Covington, TN; J.D. Schim, MD, A. Tenorio, M. Acda, The Neurology Center, Oceanside, CA; S.L. Schwartz, MD, J.S. Fischer, MD, M.S. Kipnes, MD, M.D. Blades, Diabetes and Glandular Disease Clinic, San Antonio, TX; J.S. Shymansky, MD, C. Smith, Pittsburgh Neurology Group, Pittsburgh, PA; K. Sivakumar, MD, S.A. Somers, RN, MS, CRC, Barrow Neurological Group, Phoenix, AZ; D.J. Tamulonis Jr, MD, C. Sosnowski, CNS, CNRN, St. Elizabeth Health Center, Youngstown, OH; G.L. Tan, MD, Mansfield Neurology, Mansfield, OH; R.L. Taylor, MD, K. McDonough, Taylor Medical Group, Towson, MD; D. Walk, MD, G. Parry, MD, A. Baranuskas, Y.S. Brown, University of Minnesota, Minneapolis, MN; R.M. Webb, MD, L. Neuendorf, CRC, Neurologic Associates of Tulsa, Tulsa, OK; M.I. Weintraub, MD, A.L. Rothman, MD, S. Pines-Wolert, M. Nasko, C.F. Dee, Phelps Memorial Hospital, Sleepy Hollow, NY; and G.I. Wolfe, MD, R.J. Barohn, MD, J. Ogden, RN, University of Texas, Southwestern Medical Center, Dallas, TX.

### References

- Bruyn GW, Garland H. Neuropathies of endocrine origin. In: Vinken PJ, Bruyn GW, editors. *Handbook of clinical neurology*. Vol VIII. Amsterdam: North Holland; 1970. p 29-71.
- Vinik AI. Diagnosis and management of diabetic neuropathy. *Clin Geriatr Med* 1999;15:293-316.
- Pirart J. Why don't we teach and treat diabetic patients better? *Diabetes Care* 1978;1:139-40.
- Lehtinen JM, Uusitupa M, Siitonen O, Pyorala K. Prevalence of neuropathy in newly diagnosed NIDDM and nondiabetic control subjects. *Diabetes* 1989;38:1307-13.
- Partanen J, Niskanen L, Lehtinen J, Mervaala E, Siitonen O, Uusitupa M. Natural history of peripheral neuropathy in patients with non-insulin-dependent diabetes mellitus. *N Engl J Med* 1995;333:89-94.
- Boulton AJ, Armstrong WD, Scarpello JH, Ward JD. The natural history of painful diabetic neuropathy. *Postgrad Med J* 1983;59:556-9.
- Mackel R. Properties of cutaneous afferents in diabetic neuropathy. *Brain* 1989;112:1359-76.
- Boulton AJ, Malik RA. Diabetic neuropathy. *Med Clin North Am* 1998;82:909-29.
- Baron R. Peripheral neuropathic pain. From mechanisms to symptoms. *Clin J Pain* 2000;16:S16-20.
- Kiernan MC, Hales JP, Gracies JM, Mogyoros I, Burke D. Paresthesiae induced by prolonged high-frequency stimulation of human cutaneous afferents. *J Physiol* 1997;501:461-71.
- Ochoa JL, Torebjork HE. Paresthesiae from ectopic impulse generation in human sensory nerves. *Brain* 1980;103:835-53.
- Nordin M, Nystrom B, Wallin U, Hagbarth KE. Ectopic sensory discharges and paresthesiae in patients with disorders of peripheral nerves, dorsal roots and dorsal columns. *Pain* 1984;20:231-45.
- Low PA, Dotson RM. Symptomatic treatment of painful neuropathy [editorial]. *JAMA* 1998;280:1863-4.
- Greene DA, Stevens MJ, Feldman EL. Diabetic neuropathy. Scope of the syndrome. *Am J Med* 1999;10:2S-8S.
- Kingery WS. A critical review of controlled clinical trials for peripheral neuropathic pain and complex regional pain syndromes. *Pain* 1997;73:123-39.
- Young MJ, Veves A, Breddy JL, Boulton AJ. The prediction of diabetic neuropathic foot ulceration using vibration perception thresholds. *Diabetes Care* 1994;17:557-60.
- Waxman SG. Voltage-gated ion channels in axons: localization, function and development. In: Waxman SG, Kocsis SD, Stys PK, editors. *The axon, structure, function and pathophysiology*. New York: Oxford Univ Pr; 1995. p 218-43.
- Waxman SG. Acquired channelopathies in nerve injury and MS. *Neurology* 2001;56:1621-7.
- Waxman SG. The molecular pathophysiology of pain: abnormal expression of sodium channel genes and its contributions to hyperexcitability of primary sensory neurons. *Pain* 1999;Suppl 6:S133-40.
- Horn S, Quasthoff S, Grafe P, Bostock H, Renner R, Schrank B. Abnormal axonal inward rectification in diabetic neuropathy. *Muscle Nerve* 1996;19:1268-75.
- Quasthoff S. The role of axonal ion conductances in diabetic neuropathy: a review. *Muscle Nerve* 1998;21:1246-55.
- Quasthoff S, Horn S, Grosskreutz J, Grafe P. Effects of ischemia on threshold electrotonus of peripheral nerve in diabetic patients [abstract]. *J Neurol* 1995;242:S51.
- Wall PD, Devor M. Sensory afferent impulses originate from dorsal root ganglia as well as from the periphery in normal and nerve injured rats. *Pain* 1983;17:321-7.
- Zochodne DW. Diabetic neuropathies. *Curr Treat Options Neurol* 2000;2:23-9.
- Max MB, Lynch SA, Muir J, Shoaf SE, Smoller B, Dubner R. Effects of desipramine, amitriptyline, and fluoxetine on pain in diabetic neuropathy. *N Engl J Med* 1992;326:1250-6.
- Backonja M, Beydoun A, Edwards KR, et al. Gabapentin for the symptomatic treatment of painful neuropathy in patients with diabetes mellitus. A randomized controlled trial. *JAMA* 1998;280:1831-6.
- Arner S, Meyerson BA. Lack of analgesic effects of opioids on neuropathic and idiopathic forms of pain. *Pain* 1988;33:11-23.
- Harati Y, Gooch C, Swenson M, et al. Double-blind randomized trial of tramadol for the treatment of the pain of diabetic neuropathy. *Neurology* 1998;50:1842-6.
- Low PA, Opfer-Gehrking TL, Dyck PJ, Litchy WJ, O'Brien PC. Double-blind, placebo-controlled study of the application of capsaicin cream in chronic distal painful polyneuropathy. *Pain* 1995;62:163-8.
- Treatment of painful diabetic neuropathy with topical capsaicin. A multicenter, double-blind, vehicle-controlled study. The Capsaicin Study Group. *Arch Intern Med* 1991;151:2225-9.
- Kingery WS. A critical review of controlled clinical trials for peripheral neuropathic pain and complex regional pain syndromes. *Pain* 1997;73:123-39.
- Sindrup SH, Jensen TS. Efficacy of pharmacological treatments of neuropathic pain: an update and effect related to mechanism of drug action. *Pain* 1999;83:389-400.
- Calissi PT, Jaber LA. Peripheral diabetic neuropathy: current concepts in treatment. *Ann Pharmacother* 1995;29:769-77.
- Galer BS, Ganas A, Jensen MP. Painful diabetic polyneuropathy: epidemiology, pain description and quality of life. *Diabetes Res Clin Pract* 2000;47:123-8.
- Weintraub MI. Chronic submaximal stimulation in peripheral neuropathy: is there a beneficial therapeutic relationship? Pilot study. *Am J Pain Manage* 1998;8:9-13.
- Weintraub MI. Magnetic bio-stimulation in painful diabetic peripheral neuropathy. A novel intervention—a randomized double-placebo crossover study. *Am J Pain Manage* 1999;9:8-17.
- Livingston JD. *Driving force. The natural magic of magnets*. Cambridge: Harvard Univ Pr; 1996.
- Weintraub MI. Magnetic bio-stimulation in neurologic illness. In: Weintraub MI, editor. *Alternative and complementary treatment in neurologic illness*. Philadelphia: Churchill Livingstone; 2001. p 278-86.
- Mourino MR. From Thales to Lauterbur or from the lodestone to MR imaging: magnetism and medicine. *Radiology* 1991;180:593-612.
- Geddes L. History of magnetic stimulation of the nervous system. *J Clin Neurophysiol* 1991;8:3-9.
- Macklis RM. Magnetic healing, quackery and the debate about the health effects of electromagnetic fields. *Ann Intern Med* 1993;118:376-83.
- Livingston JD. Magnetic therapy. Plausible attraction? *Skeptical Inquirer* 1998;58:25-30.
- Ramey SW. Magnetic and electromagnetic therapy. *Sci Rev Altern Med* 1998;2:13-9.
- Dyck PJ, Kratz KM, Karnes JL, et al. The prevalence by staged severity of various types of diabetic neuropathy, retinopathy, and nephropathy in a population-based cohort. The Rochester Diabetic Neuropathy Study. *Neurology* 1993;43:817-24.

45. Dyck PJ. Detection, characterization, and staging of polyneuropathy: assessed in diabetics. *Muscle Nerve* 1988;11:21-32.
46. Turk DC, Melzack R. The measurement of pain and the assessment of people experiencing pain. In: Turk DC, Melzack R, editors. *Handbook of pain assessment*. New York: Guilford Pr; 1992. p 3-14.
47. Jensen MP, Karoly P, Braver S. The measurement of clinical pain intensity. A comparison of six methods. *Pain* 1986;27:117-26.
48. Scott J, Huskisson EC. Graphic representation of pain. *Pain* 1976;2:175-84.
49. Sriwatanakul K, Kelvie W, Lasagna L, Calimlim JF, Weis OF, Mehta G. Studies with different types of visual analog scales for measurement of pain. *Clin Pharmacol Ther* 1983;34:234-9.
50. Murrin KR, Rosen M. Measurement of pain. In: Lipton S, Miles J, editors. *Persistent pain*. Vol 3. London: Academic Pr; 1983. p 17-38.
51. Masson EA, Boulton AJ. The Neurometer. Validation and comparison with conventional tests for diabetic neuropathy. *Diabetic Med* 1991;8 Spec No:S63-6.
52. Maser RE, Nielsen VK, Bass EB, et al. Measuring diabetic neuropathy: assessment and comparison of clinical examination and quantitative sensory testing. *Diabetes Care* 1989;12:270-5.
53. Arezzo JC. The use of electrophysiology for the assessment of diabetic neuropathy. *Neurosci Res Comm* 1997;21:13-23.
54. Bril V, Ellison R, Ngo M, et al. Electrophysiological monitoring in clinical trials. *Muscle Nerve* 1998;11:1368-73.
55. Serlin RC, Mendoza TR, Nakamura Y, Edwards KR, Cleland CS. When is cancer pain mild, moderate or severe? Grading pain severity by its interference with function. *Pain* 1995;61:277-84.
56. Cohen J. *Statistical power analysis for the behavioral sciences*. 2nd ed. Hillsdale: Lawrence Earlbaum Associates; 1988. p 194-5.
57. Greenhalgh T. Statistics for the non-statistician. Different types of data need different statistical tests. *BMJ* 1997;315:364-6.
58. Farrar JT, Portenoy RK, Berlin JA, Kinman JL, Strom BL. Defining the clinically important difference in pain outcome measures. *Pain* 2000;88:287-94.
59. Raja SN, Haythornthwaite JA, Pappagallo M, et al. Opioids versus antidepressants in post-herpetic neuralgia. A randomized, placebo-controlled trial. *Neurology* 2002;59:1015-21.
60. Rowbotham M, Harden N, Stacey B, Bernstein P, Magnus-Miller L. Gabapentin for the treatment of postherpetic neuralgia. A randomized, controlled trial. *JAMA* 1998;280:1837-42.
61. Segal NA, Toda Y, Huston J, et al. Two configurations of static magnetic fields for treating rheumatoid arthritis of the knee. A double-blind clinical trial. *Arch Phys Med Rehabil* 2001;82:1452-60.
62. Herrmann DN, Griffin JW, Hauer P, Cornblath DR, McArthur JC. Epidermal nerve fiber density and sural nerve morphometry in peripheral neuropathies. *Neurol* 1999;53:1634-40.
63. Waxman SG, Cummins TR, Dib-Hajj SD, Black JA. Voltage-gated sodium channels and the molecular pathogenesis of pain. A review. *J Rehabil Res Dev* 2000;37:517-28.
64. Eglon RM, Hunter JC, Dray A. Ions in the fire: recent ion-channel research and approaches to pain therapy. *Trends Pharmacol Sci* 1999;20:337-42.
65. Cavopol AV, McLean MJ, Holcomb RR. An explanatory mechanism for blockade of action potentials in neural cells by external magnetic fields [abstract]. In: *Proceedings of the Fifteenth Annual Meeting of the Bioelectromagnetics Society*; 1993 June 13-17; Los Angeles (CA). Frederick (MD): Bioelectromagnetics Society. p A-1-6.
66. Holcomb RR, Wamil AW, Pickett JD, McLean MJ. Temperature sensitivity of effects of static magnetic fields on action potentials of sensory neurons in culture [abstract]. *Soc Neurosci Abs* 1990;16:883.
67. McLean MJ, Holcomb RR, Wamil AW, Pickett JD. Effects of steady magnetic fields on action potentials of sensory neurons in vitro. *Environ Med* 1991;8:36-44.
68. McLean MJ, Holcomb RR, Wamil AW, Pickett JD, Cavopol AV. Blockade of sensory neuron action potentials by a static magnetic field in the 10 mT range. *Bioelectromagnetics* 1995;16:20-32.
69. Male J. Biological effects of magnetic fields. A possible mechanism? *Biologist* 1992;39:87-9.
70. Frankel RB, Liburdy RP. Biological effects of static magnetic fields. In: Polk C, Postow E, editors. *Handbook of biological effects of electromagnetic fields*. 2nd ed. Boca Raton: CRC Pr; 1996. p 149-83.
71. Markov MS, Colbert AP. Magnetic and electromagnetic field therapy. *J Back Musculoskeletal Rehabil* 2000;14:1-13.
72. Lednev W. Possible mechanisms for the influence of weak magnetic fields on biological systems. *Bioelectromagnetics* 1991;12:71-5.
73. Tenforde TS. Biological interactions of extremely-low-frequency electric and magnetic fields. *Bioelectrochem Bioenergetics* 1991;25:1-17.
74. Adey WR, Chopart A. Cell surface ionic phenomena in transmembrane signaling to intracellular enzyme systems. In: Blank M, Findl E, editors. *Mechanistic approaches to interactions of electromagnetic fields with living systems*. New York: Plenum Pr; 1987. p 365-87.
75. Adey WR. Tissue interactions with non-ionizing electromagnetic fields. *Physiol Rev* 1981;61:435-514.
76. Worcester DL. Structural origins of diamagnetic anisotropy in proteins. *Proc Natl Acad Sci U S A* 1978;75:5475-7.
77. Hong FT, Mauzerall D, Mauro A. Magnetic anisotropy and the orientation of retinal rods in homogeneous magnetic field. *Proc Natl Acad Sci U S A* 1971;68:1283-5.
78. Goodman R, Shirley-Henderson A. Transcription and translation in cells exposed to extremely low frequency electromagnetic fields. *Bioelectrochem Bioenergetics* 1991;25:335-55.
79. Blanchard JP, Blackman CP. Clarification and application of an ion parametric resonance model for magnetic field interactions with biological systems. *Bioelectromagnetics* 1994;15:217-38.
80. Chiabrera A, Bianco B, Catatozzolo F, et al. Electric and magnetic field effects on ligand binding to the cell membrane. In: Chiabrera A, Nicolini C, Schwan HP, editors. *Interactions between electromagnetic field and cells*. London: Plenum Pr; 1985. p 253-80.
81. McLeod BR, Liboff AR. Dynamic characteristics of membrane ions in multi-field configurations of low-frequency electromagnetic radiation. *Bioelectromagnetics* 1986;7:177-89.
82. Barker A. An introduction to the basic principles of magnetic nerve stimulation. *J Clin Neurophysiol* 1991;8:26-37.
83. Holcomb RR, Worthington WB, McCollough BA, McLean MJ. Static magnetic field therapy for pain in the abdomen and genitals. *Pediatr Neurol* 2000;23:761-4.
84. Kawakubo T, Yamauchi K, Kobayashi T. Effects of magnetic field on metabolic action in the peripheral tissue. *Jpn J Appl Physiol* 1999;38:1201-3.
85. Newrick PG, Wilson AJ, Jakubowski J, Boulton AJ, Ward JD. Sural nerve oxygen tension in diabetes. *BMJ* 1986;193:1053-4.
86. Itegin M, Gunay I, Logoglu G, Isbir T. Effects of static magnetic fields on specific adenosine-5-triphosphatase activities and bioelectrical and biomechanical properties in the rat diaphragm muscle. *Bioelectromagnetics* 1995;16:147-51.
87. Wikswow JP, Barach JP. An estimate of the steady magnetic field strength required to influence nerve conduction. *IEEE Trans Biomed Eng* 1980;27:722-3.
88. Balaban TM, Bravarenko NI, Kuznetsov AN. Influence of a stationary magnetic field on bioelectric properties of snail neurons. *Bioelectromagnetics* 1990;11:13-25.
89. Zochodne DW. The microenvironment of injured and regenerating peripheral nerves. *Muscle Nerve Suppl* 2000;9:S33-8.
90. Scarpini E, Bianchi R, Moggio M, Sciacco M, Fiori MG, Scarlato G. Decrease of nerve NA<sup>+</sup>, K(+) ATPase activity in the pathogenesis of human diabetic neuropathy. *J Neurosci* 1993;12:159-67.
91. Kennedy WR, Wendelschafer-Crabb G. The innervation of human epidermis. *J Neurol Sci* 1993;115:184-90.
92. Melzack R, Wall PD. Pain mechanisms: a new theory. *Science* 1965;150:971-8.

93. Serra J, Campero M, Ochoa J, Bostock H. Activity-dependent slowing of conduction differentiates functional subtypes of C-fibers innervating human skin. *J Physiol* 1999;515:799-811.
94. Schmidt R, Schmelz M, Forster C, Ringkamp M, Torebjork E, Handwerker H. Novel classes of responsive and unresponsive C nociceptors in human skin. *J Neurol Sci* 1995;15:333-41.
95. Cummins TR, Waxman SG. Downregulation of tetrodotoxin-resistant sodium currents and upregulation of a rapidly repriming tetrodotoxin-sensitive sodium current in small spinal sensory neurons after nerve injury. *J Neurosci* 1997;1:3503-14.
96. Waling K, Sundelin G, Ahlgren C, Jarvholm B. Perceived pain before and after three exercise programs—a controlled clinical trial of women with work-related trapezius myalgia. *Pain* 2000;85:201-7.
97. Raymond I, Nielsen TA, Lavigne G, Manzini C, Choiniere M. Quality of sleep and its daily relationship to pain intensity in hospitalized adult burn patients. *Pain* 2001;92:381-8.
98. Von Korff M, Ormel J, Keefe FJ, Dworkin SF. Grading the severity of chronic pain. *Pain* 1992;50:133-49.
99. Moscucci M, Byrne L, Weintraub M, Cox C. Blinding, unblinding, and the placebo effect: an analysis of patients' guesses of treatment assignment in a double-blind clinical trial. *Clin Pharmacol Ther* 1987;41:259-65.
100. Noseworthy JH, Ebers GC, Vandervoort MK, Farquhar RE, Yetisir E, Roberts R. The impact of blinding on the results of a randomized, placebo-controlled multiple sclerosis clinical trial. *Neurology* 1994;44:16-20.
101. Periquet MI, Novak V, Collins MP, et al. Painful sensory neuropathy: prospective evaluation of painful feet using electrodiagnosis and skin biopsy. *Neurology* 1999;53:1641-7.
102. Lauria G, McArthur JC, Hauer PE, Griffin JW, Cornblath DR. Neuropathologic alterations in diabetic truncal neuropathy. Evaluation by skin biopsy. *J Neurol Neurosurg Psychiatry* 1998;65:762-6.
103. McArthur JC, Yiannoutsos C, Simpson DM, et al. A phase II trial of nerve growth factor for sensory neuropathy associated with HIV infection. AIDS Clinical Trials Group Team 291. *Neurology* 2000;54:1080-8.

#### Suppliers

- a. MMT-6J04-VH; Magnet-Physics Inc, 770 W Algonquin Rd, Arlington Heights, IL 60005.
- b. Nu-Magnetics Inc, 6 N Wind Dr, Port Jefferson, NY 11777.
- c. Nikken Inc, 52 Discovery, Irvine, CA 92618.
- d. Neurotron Inc, 1501 Sulgrave Ave, Ste 203, Baltimore, MD 21209.
- e. Version 10.0; SPSS Inc, 233 S Wacker Dr, Chicago, IL 60606.